arXiv:1612.00932v1 [math.FA] 3 Dec 2016

QUATERNIONIC FOCK SPACE IN AN EUCLIDEAN UNIT BALL $\mathbb{B}^n$

SANJAY KUMAR AND KHALID MANZOOR

Abstract. In this paper, we study the quaternionic counterpart of complex Fock spaces $\mathfrak{F}^p_\alpha (0 < p < \infty$ and for some parameter $\alpha)$ of entire slice hyperholomorphic functions in an Euclidean unit ball $\mathbb{B}^n$ in $\mathbb{H}^n$. We also study some basic properties of these spaces.

1. Introduction

Consider the 4-dimensional real algebra $\mathbb{H} = \{x_0 + x_1 i + x_2 j + x_3 k : x_l \in \mathbb{R}$ for $0 \leq l \leq 3\}$ of quaternions. Then $\{1, i, j, k\}$ form the standard basis of quaternions with imaginary units, where $Re(q) = x_0$, $Im(q) = x_1 i + x_2 j + x_3 k$ denote the real part and imaginary part of q respectively. The conjugate of q is given by $\bar{q} = Re(q) - Im(q) = x_0 - (x_1 i + x_2 j + x_3 k)$. By $|q| = \sqrt{q\bar{q}} = \sqrt{\sum_{l=0}^{3} x_l^2}$, for $x_l \in \mathbb{R}$, we mean the Euclidean norm on $\mathbb{H}$. For any non-zero quarternion q, its multiplicative inverse is given by $\frac{\bar{q}}{|q|^2}$. The set $\mathbb{S} = \{q \in \mathbb{H} : q = x_1 i + x_2 j + x_3 k$ and ${x_1}^2 + {x_2}^2 + {x_3}^2 = 1\}$ denotes the two-dimensional unit sphere of purely imaginary quaternions. Any element $I \in \mathbb{S}$ is such that $I^2 = -1$. This implies that the elements of S are imaginary units. The quaternion is considered as the union of complex plane $\mathbb{C}_I = \mathbb{R} + \mathbb{R}I$ (also called slices), each one is identified by an imaginary unit $I \in \mathbb{S}$. Let $\Omega_I = \Omega \cap \mathbb{C}_I$, for some domain Ω of $\mathbb{H}$. For any quaternion q we can write $q = x_0 + x_1 i + x_2 j + x_3 k = x_0 + Im(q) = x_0 + |Im(q)| I_q = x + y I_q$, with $I_q = \frac{Im(q)}{|Im(q)|}$ if $|Im(q)| \neq 0$, otherwise we take arbitrary I in $\mathbb{S}$.

For some more details on slice holomorphic functions one can refer to [18, 19] and references therein. In recent times the notion of slice holomorphic functions has been studied intensively. The Hardy spaces, Bloch spaces, Dirichlet spaces, etc. are studied in the slice hyperholomorphic settings. For Fock spaces in the slice hyperholomorphic settings, see [10].

For details about recent work on slice hyperholomorphic and their applications to Operator Theory, Schur Analysis, Quantum Physics, etc one can refer to an excellent

Date: December 6, 2016.

2000 *Mathematics Subject Classification.* Primary 47B38, 47B33, 30D55.

Key words and phrases. Fock space, slice regular Fock space, slice hyperholomorphic functions.

recent survey [3, 7, 18, 19, 25] and references therein. For study of Fock spaces of holomorphic functions, we can refer to [32].

The work of this paper is motivated by [10, 29] and recent work of Ueki [27, 28]. Here, we begin with some basic results in the quaternionic-valued slice regular functions.

Definition 1.1. *Let Ω be a domain in $\mathbb{H}$. A real differentiable function $f : \Omega \to \mathbb{H}$ is said to be the (left) slice regular or slice hyperholomorphic if for any $I \in \mathbb{S}, f_I$ is holomorphic in Ω_I, i.e.,*

$$\left(\frac{\partial}{\partial x} + I\frac{\partial}{\partial y}\right) f_I(x + yI) = 0,$$

where f_I denote the restriction of f to Ω_I. The class of slice regular functions on Ω is denoted by $SR(\Omega)$.

For slice regular functions, we have the following useful result.

Theorem 1.2. [22, Theorem 2.7] *A function $f : \mathbb{B} \to \mathbb{H}$ is said to be slice regular if and only if it has a power series of the form*

$$f(q) = \sum_{n=0}^{\infty} q^n a_n, \text{ where } a_n = \frac{1}{n!}\frac{\partial^n f(0)}{\partial x^n} \tag{1}$$

converging uniformly on $\mathbb{B}$.

Lemma 1.1. [18, Lemma 4.1.7]*(Spliting Lemma) If f is a slice regular function on the domain Ω, then for any $I, J \in \mathbb{S}$, with $I \perp J$ there exist two holomorphic functions $F, L : \Omega_I = \Omega \cap \mathbb{C}_I \to \mathbb{C}_I$ such that*

$$f_I(z) = F(z) + L(z)J; \text{ for any } \ z = x + yI \in \Omega_I. \tag{2}$$

One of the most important property of the slice regular functions is their Representation Formula. It only holds on the open sets which are stated below.

Definition 1.3. *Let Ω be an open set in $\mathbb{H}$. We say Ω is axially symmetric if for any $q = x + yI_q \in \Omega$, all the elements $x + yI$ are contained in Ω, for all $I \in \mathbb{S}$ and Ω is said to be slice domain if $\Omega \cap \mathbb{R}$ is non empty and $\Omega \cap \mathbb{C}_I$ is a domain in $\mathbb{C}_I$ for all $I \in \mathbb{S}$.*

Theorem 1.4. [18, Theorem 4.3.2] *(Representation Formula) Let f be a slice regular function in the domain $\Omega \subset \mathbb{H}$. Then for any $J \in \mathbb{S}$ and for all $z = x + yI \in \Omega$ such that*

$$f(x + yI) = \frac{1}{2}\{(1 + IJ)f(x - yI) + (1 - IJ)f(x + yI)\}.$$

Remark 1.5. *Let I, J be orthogonal imaginary units in $\mathbb{S}$ and Ω be an axially symmetric slice domain. Then the Splitting Lemma and the Representation formula generate a class of operators on the slice regular functions as follows:*

$$Q_I : SR(\Omega) \to hol(\Omega_I) + hol(\Omega_I)J$$
$$Q_I : f \mapsto f_1 + f_2 J$$

$$P_I : hol(\Omega_I) + hol(\Omega_I)J \to SR(\Omega)$$
$$P_I[f](q) = P_I[f](x + yI_q) = \frac{1}{2}[(1 - II_q)f(x + yI) + (1 + II_q)f(x - yI)],$$

where $hol(\Omega_I)$ *is the space of all holomorphic functions on* Ω_I *for some* $I \in \mathbb{S}$. *Also,*

$$P_I \circ Q_I = \mathrm{I}_{SR(\Omega)} \text{ and } Q_I \circ P_I = \mathrm{I}_{SR(hol(\Omega_I)+hol(\Omega_I))},$$

where I *is an identity operator.*

Since pointwise product of functions does not preserve slice regularity, a new multiplication operation for regular functions is defined. In the special case of power series, the regular product (or $\star-$product) of $f(q) = \sum_{n=0}^{\infty} q^n a_n$ and $g(q) = \sum_{n=0}^{\infty} q^n b_n$ is

$$f \star g(q) = \sum_{n \geq 0} q^n \sum_{k=0}^{n} a_k b_{n-k}.$$

The $\star-$product is related to the standard pointwise product by the following formula.

Theorem 1.6. [12, Proposition 2.4] *Let* f, g *be regular functions on* $\mathbb{B}$. *Then* $f \star g(q) = 0$ *if* $f(q) = 0$ *and* $f(q)g(f(q)^{-1}qf(q))$ *if* $f(q) \neq 0$. *The reciprocal* $f^{-\star}$ *of a regular function* $f(q) = \sum_{n=0}^{\infty} q^n a_n$ *with respect to the* $\star-$*product is*

$$f^{\star}(q) = \frac{1}{f \star f^c(q)} f^c(q),$$

where $f^c(q) = \sum_{n=0}^{\infty} q^n \overline{a_n}$ *is the regular conjugate of* f. *The function* $f^{-\star}$ *is regular on* $\mathbb{B} \setminus (q \in \mathbb{B} | f \star f^c(q) = 0)$ *and* $f \star f^{-\star} = 1$ *there.*

2. Fock space

Let $\mathbb{C}^n$ be n-dimensional Euclidean space and dA be the normalized area measure on $\mathbb{C}^n$. For $\alpha > 0$ and $0 < p < \infty$, a holomorphic function $f : \mathbb{C}^n \to \mathbb{C}$ is said to be in Fock space $\mathfrak{F}_{p,\mathbb{C}}$ if

$$\left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{C}^n} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA(z) < \infty,$$

where $z = (z_1, z_2 ..... , z_n)$ in $\mathbb{C}^n$, $dA_n(z) = (\frac{1}{\pi})^n \prod_{k=1}^{n} dx_k dy_k,\ z_k = x_k + j y_k,\ x_k, y_k \in \mathbb{R}$. For $z = (z_1, z_2, ... , z_n)$ and $w = (w_1, w_2, ... , w_n)$ in $\mathbb{C}^n$, we write $\langle z, w \rangle_\alpha$ for the Euclidean inner product $\sum_{k=1}^{n} z_k \overline{w_k}$ and $|z| = \sqrt{\langle z, z \rangle}$. Let $\mathbb{B}^n(0,1) = \mathbb{B}^n = \{q = x + yI_q = (q_1, q_2, ... , q_n) : \sum_{k=1}^{n} |q_k|^2 < 1\}$ be the quaternionic unit ball centered at origin in $\mathbb{H}^n$ and $\mathbb{B}^n \cap \mathbb{C}_I^n = \mathbb{B}_I^n$ denote n-dimensional unit disk in the complex plane $\mathbb{C}_I^n$ for $I \in \mathbb{S}$. Now we begin with the following definitions.

Definition 2.1. *For $0 < p < \infty, \alpha > 0$ and some $I \in \mathbb{S}$, the quaternionic right linear space of entire slice regular functions f is said to be the quaternionic slice regular Fock space on the unit ball $\mathbb{B}^n$, if for any $q = x + yI_q \in \mathbb{B}^n$*

$$\left(\frac{\alpha}{\pi}\right)^n \sup_{I\in\mathbb{S}} \int_{\mathbb{B}_I^n} \left| f(q) e^{\frac{-\alpha}{2}|q|^2} \right|^p dA_I(q) < \infty,$$

that is,

$$\mathfrak{F}_\alpha^p = \left\{ f \in SR(\mathbb{B}^n, \mathbb{H}^n) : \left(\frac{\alpha}{\pi}\right)^n \sup_{I\in\mathbb{S}} \int_{\mathbb{B}_I^n} \left| f(q) e^{\frac{-\alpha}{2}|q|^2} \right|^p dA_I(q) \right\} < \infty,$$

where $dA_I(q) = \frac{1}{\pi} dxdy$ and is Möbius invariant measure on $\mathbb{B}^n$. If we write

$$d\lambda_{\alpha,I}(q) = \left(\frac{\alpha}{\pi}\right)^n e^{-\alpha|q|^2} dA_I(q),$$

then $\mathfrak{F}_\alpha^2$ is a Hilbert space (complete inner product space) with inner product $\langle ., . \rangle_\alpha$ defined as

$$\langle f, g \rangle_\alpha = \int_{\mathbb{B}_I^n} f(q) \overline{g(q)} d\lambda_{\alpha,I}(q)$$

with the norm given by

$$\|f\|_{\mathfrak{F}_\alpha^p} = \left(\frac{\alpha}{\pi}\right)^{np} \sup_{I\in\mathbb{S}} \left(\int_{\mathbb{B}_I^n} \left| f(q) e^{\frac{-\alpha}{2}|q|^2} \right|^p dA_I(q) \right)^{\frac{1}{p}}.$$

By $\mathfrak{F}_{\alpha,I}^p$, we denote the quaternionic right linear space of entire slice regular functions on $\mathbb{B}^n$ such that

$$\left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}_I^n} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) < \infty.$$

Furthermore, for each function $f \in \mathfrak{F}_{\alpha,I}^p$, we define

$$\|f\|_{\mathfrak{F}_{\alpha,I}^p} = \left(\frac{\alpha}{\pi}\right)^{np} \left(\int_{\mathbb{B}_I^n} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) : z = x + yI \in \mathbb{B}^n \cap \mathbb{C}_I^n \right)^{\frac{1}{p}}.$$

Remark 2.2. [29, P. 499] *Let $I \in \mathbb{S}$ be such that $J \perp I$. Then there exist holomorphic functions $f_1, f_2 : \mathbb{B}_I^n \to \mathbb{C}_I$ such that $Q_I[f] = f_1 + f_2 J$ for some holomorphic map $Q_I[f]$ in complex variable $z = (z_1, z_2, ...z_n) \in \mathbb{B}_I^n$. Then, we have*

$$\begin{aligned} \left| f_l(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p &\leq \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p \\ &\leq 2^{max\{0,p-1\}} \left| f_1(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p + 2^{max\{0,p-1\}} \left| f_2(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p. \end{aligned}$$

The condition $f \in \mathfrak{F}_{\alpha,I}^p$ is equivalent to f_1 and f_2 belonging to n-dimensional complex Fock space.

We can easily prove the following result.

Proposition 2.3. *Let $I \in \mathbb{S}$ and $\alpha > 0$. Then $f \in \mathfrak{F}^p_{\alpha,I}, p > 1$ if and only if $f \in \mathfrak{F}^p_\alpha$. Moreover, the spaces $(\mathfrak{F}^p_{\alpha,I}, \|.\|_{\mathfrak{F}^p_{\alpha,I}})$ and $(\mathfrak{F}^p_\alpha, \|.\|_{\mathfrak{F}^p_\alpha})$ have equivalent norms. More precisely, one has*

$$\|f\|^p_{\mathfrak{F}^p_{\alpha,I}} \leq \|f\|^p_{\mathfrak{F}^p_\alpha} \leq 2^p \|f\|^p_{\mathfrak{F}^p_{\alpha,I}}.$$

Proposition 2.4. *Let $I, J \in \mathbb{S}$ and let $p > 1, \alpha > 0$ and $f \in SR(\mathbb{B}^n, \mathbb{H}^n)$. Then $f \in \mathfrak{F}^p_{\alpha,I}$ if and only if $f \in \mathfrak{F}^p_{\alpha,J}$.*

Proof. Since $f \in SR(\mathbb{B}^n, \mathbb{H}^n)$ and $w = x + yJ \in \mathbb{B}^n_J$ and $z = x + yI \in \mathbb{B}^n_I$, with $|w| = |z|$. By Representation Formula, we have

$$f(w) = f(x + yJ) = \frac{1}{2} |(1 - JI)f(z) + (1 + JI)f(\bar{z})| \leq |f(z)| + |f(\bar{z})|.$$

This implies that

$$\begin{aligned} \int_{\mathbb{B}^n_J} \left| f(w) e^{\frac{-\alpha}{2}|w|^2} \right|^p dA_J(w) &\leq 2^{max\{p-1,0\}} \int_{\mathbb{B}^n_I} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) \\ &+ 2^{max\{p-1,0\}} \int_{\mathbb{B}^n_I} \left| f(\bar{z}) e^{\frac{-\alpha}{2}|\bar{z}|^2} \right|^p dA_I(\bar{z}). \end{aligned}$$

As $\bar{z} \to z$, we have

$$\left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_J} \left| f(w) e^{\frac{-\alpha}{2}|w|^2} \right|^p dA_J(w) \leq 2^{max\{p,1\}} \left(\left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_I} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) \right).$$

Thus, for any $f \in \mathfrak{F}^p_{\alpha,I}$, we see $f \in \mathfrak{F}^p_{\alpha,J}$. By interchanging the roles of I and J, we obtain other result. ∎

Proposition 2.5. *Suppose $p > 1$ and $\alpha > 0$. If f is in $SR(\mathbb{B}^n, \mathbb{H}^n)$, then following assertations are equivalent:*

(a) $f \in \mathfrak{F}^p_\alpha(\mathbb{B}^n)$;
(b) $f \in \mathfrak{F}^p_{\alpha,I}(\mathbb{B}^n_I)$ *for some* $I \in \mathbb{S}$.

Proof. To prove this, it is sufficient to show $(b) \Rightarrow (a)$. Suppose $f \in \mathfrak{F}^p_{\alpha,I}$. Let $q = x + yI_q$ and $z = x + yI$. Applying Representation Formula followed by triangle inequality and the fact that $|q| = |z| = |\bar{z}|$, we see that

$$f(q) = \frac{1}{2} |(1 - I_q I)f(x + yI) + (1 + I_q I)f(x - yI)| \leq |f(z)| + |f(\bar{z})|.$$

This implies that

$$\begin{aligned} \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_I} \left| f(q) e^{\frac{-\alpha}{2}|q|^2} \right|^p dA_I(q) &\leq 2^{max\{p-1,0\}} \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_I} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) \\ &+ 2^{max\{p-1,0\}} \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_I} \left| f(\bar{z}) e^{\frac{-\alpha}{2}|\bar{z}|^2} \right|^p dA_I(\bar{z}) \\ &\leq 2^{max\{p,1\}} \left(\frac{\alpha}{\pi}\right)^n \int_{\mathbb{B}^n_I} \left| f(z) e^{\frac{-\alpha}{2}|z|^2} \right|^p dA_I(z) \\ &< \infty. \end{aligned}$$

Thus, the condition (a) holds. ∎

Remark 2.6. *By $L^p(\mathbb{B}_I, d\lambda_{\alpha,I}, \mathbb{H})$ we define the set of functions $g : \mathbb{B}_I^n \to \mathbb{H}$ such that*

$$\int_{\mathbb{B}_I^n} |g(w)|^p d\lambda_{\alpha,I}(w) < \infty,$$

where $d\lambda_{\alpha,I}(w) = \frac{\alpha}{\pi} e^{-\alpha|z|^2} dA_I(w)$, for $\alpha > 0$ is called the Gaussian probability measure. Note that for $J \in \mathbb{S}$ with $J \perp I$ and $g = g_1 + g_2 J$ with $g_1, g_2 : \mathbb{B}_I^n \to \mathbb{C}_I$, then $g \in L^p(\mathbb{B}_I, d\lambda_{\alpha,I}, \mathbb{H})$ if and only if $g_1, g_2 \in L^p(\mathbb{B}_I, d\lambda_{\alpha,I}, \mathbb{C}_I)$.
Clearly, $\mathfrak{F}_\alpha^p$ is closed subspace of $L^p(\mathbb{B}_I, d\lambda_{\alpha,I}, \mathbb{H})$. In complex analysis, the reproducing kernel of complex Fock space for $p = 2$ is given by

$$K_\alpha^{\mathbb{C}_I^n}(z, w) = e^{\alpha\langle z, w\rangle};\ z, w \in \mathbb{C}_I^n.$$

This gives the motivation for the following definition.

Definition 2.7. *For any $q \in \mathbb{B}^n$, the slice regular exponential function is given by*

$$e^q = \sum_{n=0}^{\infty} \frac{q^n}{n!}.$$

Let $e^{zw} = \sum_{n=0}^{\infty} \frac{z^n w^n}{n}$ be a holomorphic function in variable z in the complex plane $\mathbb{C}_I^n$. Clearly, e^{zw} is not slice regular in both variable. Setting $e_\star^{qw} = \sum_{n=0}^{\infty} \frac{q^n w^n}{n!}$, then we see that the function is left slice regular in q and right slice regular in w, where $\star$ denote the slice regular product. By Representation formula, we can obtain the extension of function e^{zw} to $\mathbb{H}$, as

$$ext(e^{zw}) = \frac{1}{2}\{(1 - IJ)e^{zw} + (1 + IJ)e^{\bar{z}w}\} = e^{qw},$$

where $q \in \mathbb{B}^n$ and for some arbitrary w. For $I_q = \frac{Im(q)}{|Im(q)|} \in \mathbb{S}$ and $\alpha > 0$, we define

$$B_\alpha(q, w) = e_\star^{\alpha q \bar{w}} \ for\ each\ q = x + yI_q \in \mathbb{B}^n,$$

and is called slice regular reproducing kernel of quaternionic Fock space.

The following theorem is a quaternionic version of [30, Theorem 3.1].

Theorem 2.8. *Let $I \in \mathbb{S}$. Then for any $p > 1$, there exists a sequence $\{z_k\}_{k>1}$ in $\mathbb{B}_I^n$ with the following property: An entire function $f \in \mathfrak{F}_\alpha^p$ if and only if*

$$f(z) = P_I \sum_{k=1}^{\infty} w_{z_k}(z) a_k, \tag{3}$$

where $\{a_k\} \in l^p(\mathbb{H}^n)$ and $w_{z_k}(z) = e_\star^{\alpha z \bar{z_k} - \frac{\alpha}{2}|z_k|^2}$ is the slice regular normalized reproducing kernel with $P_I = \frac{\vec{z}}{\|\vec{z}\|}$.

Proof. Let $J \in \mathbb{S}$ be such that $J \perp I$. Let $f \in \mathfrak{F}^p_\alpha$. Then $f \in \mathfrak{F}^p_{\alpha,I}$ for $I \in \mathbb{S}$. If we restrict f on $\mathbb{B}^n_I$, then we have $f_I(z) = Q_I[f](z) = f_1(z) + f_2(z)J$ with holomorphic functions $f_1, f_2 : \mathbb{B}^n \cap \mathbb{C}^n_I \to \mathbb{C}_I$. Applying [30, Theorem 3.1] to f_1 and f_2, we can find the sequences $\{a_{1,k}\}_{k\geq 1}$ and $\{a_{2,k}\}_{k\geq 1}$ in $l^p(\mathbb{C}^n_I)$ such that

$$f_1(z) = \sum_{k=1}^{\infty} w_{z_k}(z)a_{1,k} \text{ and } f_2(z) = \sum_{k=1}^{\infty} w_{z_k}(z)a_{2,k}.$$

So, we can write

$$Q_I[f](z) = f_1(z) + f_2(z)J = \sum_{k=1}^{\infty} w_{z_k}(z)a_{1,k} + \sum_{k=1}^{\infty} w_{z_k}(z)a_{2,k}J.$$

As $P_I \circ Q_I[f] = \mathrm{I}_{SR(\mathbb{B}^n_I)}$ an identity operator, we have

$$\begin{aligned} f = P_I \circ Q_I[f](z) &= P_I \left\{ \sum_{k=1}^{\infty} w_{z_k}(z)a_{1,k} + \sum_{k=1}^{\infty} w_{z_k}(z)a_{2,k}J \right\} \\ &= P_I \left\{ \sum_{k=1}^{\infty} w_{z_k}(z) \left(a_{1,k} + a_{2,k}J\right) \right\} \\ &= P_I \left\{ \sum_{k=1}^{\infty} w_{z_k}(z)a_k \right\}, \end{aligned}$$

where $a_k = a_{1,k} + a_{2,k}J \in l^p(\mathbb{H}^n)$. Conversely, suppose the condition (3) holds. We claim $f \in \mathfrak{F}^p_\alpha$. Then for any $J \in \mathbb{S}$, we have $a_k = a_{1,k} + a_{2,k}J$, where $\{a_{l,k}\}_{k\geq 1} \in l^p(\mathbb{C}^n_I),\ l = 1, 2$. So, we can write

$$Q_I[f](z) = \sum_{k=1}^{\infty} a_{1,k}w_{z_k}(z) + \sum_{k=1}^{\infty} a_{2,k}w_{z_k}(z)J.$$

Since $\{a_k\}_{k\geq 1}$ lie in $l^p(\mathbb{H}^n)$, it follows that $\{a_{l,k}\}_{k\geq 1}$ belong to $l^p(\mathbb{C}^n_I)$. This implies that f_1, f_2 lie in the complex Fock space $\mathfrak{F}^p_{\alpha,I,\mathbb{C}_I}$ which is equivalent to $f \in \mathfrak{F}^p_{\alpha,I}$ and hence $f \in \mathfrak{F}^p_\alpha$. ∎

The proof of the following proposition is analogus to [29, Theorem 2.19].

Theorem 2.9. *Suppose $1 \leq p < \infty$ and $\alpha > 0$. If $f \in \mathfrak{F}^p_\alpha$ with some $I \in \mathbb{S}$, then for any $z \in \mathbb{B}^n_I$, there exists some $M > 0$ such that*

$$|f(z)| \leq 2^{max\{p,1\}} M \|f\|_{\mathfrak{F}^p_\alpha}.$$

Proposition 2.10. *Let $1 < p < \infty$ and $\alpha > 0$. If $\{f_n\}_{n\in\mathbb{N}}$ is any sequence in $\mathfrak{F}^p_\alpha(\mathbb{B}^n)$, then $\{f_n\}$ converges weakly to $f \in \mathfrak{F}^p_\alpha(\mathbb{B}^n)$ if and only if there exists $\lambda > 0$ such that $\|f_n\|_{\mathfrak{F}^p_\alpha} \leq \lambda$ and $f_n \to f$ uniformly on the compacts subsets of $\mathbb{B}^n$.*

Proof. Suppose sequence f_n is weakly convergent in $\mathfrak{F}^p_\alpha$ and so in $\mathfrak{F}^p_{\alpha,I}$. Let $J \in \mathbb{S}$ be such that $J \perp I$ and let $f_{n,1}, f_{n,2}$ be holomorphic functions such that $Q_I[f_n] = f_{n,1} + f_{n,2}J$. By Remark 2.2, $f_{n,1}, f_{n,2}$ lie in the complex Fock space. Then, from

corresponding result in the complex case, we see that the sequences $\{f_{n,1}\}$ and $\{f_{n,2}\}$ converges weakly to functions f_1 and f_2 as $n \to \infty$, respectively if and only if there exists some positive constant M such that $\|f_{n,l}\|_{\mathfrak{F}^p_{\alpha,\mathbb{C}_I}} \leq M$ and $f_{n,l} \to f_l$ uniformly on the compact subsets of $\mathbb{B}^n_I$, for $l = 1, 2$ if and only if $\|f_n\|_{\mathfrak{F}^p_{\alpha,I}} \leq 2^{p-1}\|f_{n,l}\|_{\mathfrak{F}^p_{\alpha,\mathbb{C}_I}} \leq 2^{p-1}M = \lambda$ for some $\lambda = 2^{p-1}M$ and for any $f = p_I[f_1 + f_2 J]$ in $\mathfrak{F}^p_{\alpha,I}$,

$$|f_n - f| \leq |f_{n,1} - f_1| + |f_{n,2} - f_2| \to 0, \text{ as } n \to \infty$$

uniformly on the compacts subsets of $\mathbb{B}^n_I$. ■

Now, we have the following definitions.

Definition 2.11. *By space $\mathfrak{F}_{\infty,\alpha}$, we mean the space of all slice regular Fock functions f such that*

$$\sup_{q\in\mathbb{B}^n} |f(q)|e^{\frac{-\alpha}{2}|q|^2} < \infty,$$

that is,

$$\mathfrak{F}_{\infty,\alpha} = \left\{ f \in SR(\mathbb{B}^n, \mathbb{H}^n) : \sup |f(q)|e^{\frac{-\alpha}{2}|q|^2} : \ q = x + yI_q \in \mathbb{B}^n \right\} < \infty.$$

The space $\mathfrak{F}_{\infty,\alpha}$ is a Banach space under the norm defined as

$$\|f\|_{\mathfrak{F}_{\infty,\alpha}} = \sup \left\{ |f(q)|e^{\frac{-\alpha}{2}|q|^2} : \ q \in \mathbb{B}^n \right\}.$$

Definition 2.12. *We define the space $\mathfrak{F}_{\infty,\alpha,I}$ as the quaternionic right linear space of all slice regular functions f defined on $\mathbb{B}^n_I$ such that*

$$\|f\|_{\mathfrak{F}_{\infty,\alpha,I}} = \sup \left\{ |Q_I[f](z)|e^{\frac{-\alpha}{2}|z|^2} : \ z = x + yI \in \mathbb{B}^n_I \right\} < \infty.$$

Definition 2.13. *The space $\mathfrak{F}^0_{\infty,\alpha}$ is a closed subspace of $\mathfrak{F}_{\infty,\alpha}$ and is defined as the space of all slice regular functions in $\mathbb{B}^n$ such that*

$$\lim_{|q|\to\infty} |f(q)|\, e^{\frac{-\alpha}{2}|q|^2} = 0.$$

Remark 2.14. *Let $I, J \in \mathbb{S}$ be such that $I \perp J$ and let $f \in SR(\mathbb{B}^n, \mathbb{H}^n)$. Furthermore, let $f_1, f_2 : \mathbb{B}^n_I \to \mathbb{C}_I$ be holomorphic functions in the complex Fock space $\mathfrak{F}^0_{\infty,\alpha,\mathbb{C}_I}$ such that $Q_I[f] = f_1 + f_2 J$. Then*

$$\left| f(z)e^{\frac{-\alpha}{2}|z|^2} \right|^2 = \left| f_1(z)e^{\frac{-\alpha}{2}|z|^2} \right|^2 + \left| f_2(z)e^{\frac{-\alpha}{2}|z|^2} \right|^2 \text{ for all } z \in \mathbb{B}^n_I.$$

By Representation Formula, it follows that $f \in \mathfrak{F}^0_{\infty,\alpha}$ if and only if f_1, f_2 lie in the complex space $\mathfrak{F}^0_{\infty,\alpha,\mathbb{C}_I}$.

By Representation Formula, we have the following result.

Proposition 2.15. *Suppose $I \in \mathbb{S}$ and $\alpha > 0$. Then $f \in \mathfrak{F}_{\infty,\alpha,I}$ if and only if $f \in \mathfrak{F}_{\infty,\alpha}$. Futhermore, the spaces $(\mathfrak{F}_{\infty,\alpha,I}, \ \|.\|_{\mathfrak{F}_{\infty,\alpha,I}})$ and $(\mathfrak{F}_{\infty,\alpha}, \ \|.\|_{\mathfrak{F}_{\infty,\alpha}})$ have equivalent norms. More precisely, one has*

$$\|f\|_{\mathfrak{F}_{\infty,\alpha,I}} \leq \|f\|_{\mathfrak{F}_{\infty,\alpha}} \leq 2\|f\|_{\mathfrak{F}_{\infty,\alpha,I}}.$$

Proposition 2.16. *For $p > 1$ and $\alpha > 0$ with multi-index m such that $m = (m_1, m_2, ..., m_n)$ and $|m| = |m_1| + |m_2| + ... + |m_n|$. Then for all $f \in SR(\mathbb{B}^n, \mathbb{H}^n)$ and all $z \in \mathbb{B}_I^n$, for some $I \in \mathbb{S}$*

$$\|a_m z^m\|_{\mathfrak{F}_{\infty,\alpha}} \leq 2^{max\{p,1\}} \prod_k \sqrt{\frac{m_k}{2}} \sup_{z \in \mathbb{B}_I^n} |f(z)| e^{\frac{-\alpha}{2}|z|^2}.$$

Proof. Let $J \in \mathbb{S}$ with $I \perp J$. Then by Splitting Lemma, we can find two holomorphic functions $f_1, f_2 \in hol(\mathbb{B}_I^n)$ such that $Q_I[f] = f_1 + f_2 J$. Therefore from [28, Lemma 2], we conclude that

$$\begin{aligned}
\|a_m z^m\|_{\mathfrak{F}_{\infty,\alpha}} &\leq 2^{max\{p-1,0\}} \left(\|a_{m,1} z^m\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}} + \|a_{m,2} z^m\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}} \right) \\
&\leq 2^{max\{p-1,0\}} \left(\prod_k \sqrt{\frac{m_k}{2}} \sup_{z \in \mathbb{B}_I^n} |f_1(z)| e^{\frac{-\alpha}{2}|z|^2} + \prod_k \sqrt{\frac{m_k}{2}} \sup_{z \in \mathbb{B}_I^n} |f_2(z)| e^{\frac{-\alpha}{2}|z|^2} \right) \\
&\leq 2^{max\{p-1,0\}} \prod_k \sqrt{\frac{m_k}{2}} \sup_{z \in \mathbb{B}_I^n} (|f_1(z)| + |f_2(z)|) e^{\frac{-\alpha}{2}|z|^2} \\
&\leq 2^{max\{p,1\}} \prod_k \sqrt{\frac{m_k}{2}} \sup_{z \in \mathbb{B}_I^n} (|f(z)|) e^{\frac{-\alpha}{2}|z|^2}.
\end{aligned}$$

Hence the result. ∎

Proposition 2.17. *For $0 < r < 1$, the space $\mathfrak{F}_{\infty,\alpha}$ is the closure with respect to norm $\|.\|_{\mathfrak{F}_{\infty,\alpha}}$ of the set of quaternionic polynomials of the form*

$$P_k^{(r)}(q) = \sum_{l=1}^{k} \sum_{|m|=l} r^{|m|} q^m a_m \ for \ all \ a_m \in \mathbb{H}^n.$$

In particular, $\mathfrak{F}_{\infty,\alpha}$ is separable.

Proof. Let $f \in \mathfrak{F}_{\infty,\alpha}$. Then by orthogonality of imaginary units I, J one can write $f_I = Q_I[f] = f_1 + f_2 J$, where f_1, f_2 are holomorphic functions in the complex Fock space. Since the polynomials are dense in the complex Fock space, from [28, Lemma 3], we find the complex polynomials of the form

$$P_{k,1}^{(r)}(z) = \sum_{l=1}^{k} \sum_{|m|=l} r^{|m|} z^m \beta_{k,m} \text{ and } P_{k,2}^{(r)}(z) = \sum_{l=1}^{k} \sum_{|m|=l} r^{|m|} z^m \gamma_{k,m}$$

such that

$$\|f_n - P_{k,n}^{(r)}\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I^n}} \to 0, \quad \text{as } k \to \infty \text{ for } n = 1, 2.$$

Now, let $a_{k,m} = \beta_{k,m} + \gamma_{k,m} J$. Then, we have

$$P_k^{(r)}(q) = \sum_{l=1}^{k} \sum_{|m|=l} r^{|m|} q^m \left(\beta_{k,m} + \gamma_{k,m} J\right) = \sum_{l=1}^{k} \sum_{|m|=l} r^{|m|} q^m a_{k,m}.$$

Therefore, by Remark 2.15, it follows that

$$\begin{aligned}\|f-P_k^{(r)}\|_{\mathfrak{F}_{\infty,\alpha}} &\leq 2\|f-P_k^{(r)}\|_{\mathfrak{F}_{\infty,\alpha,I}}\\ &\leq 2\|f_1-P_{k,1}^{(r)}\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}}+\|f_2-P_{k,2}^{(r)}\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}}\\ &\to 0,\ as\ k\to\infty.\end{aligned}$$

Hence the space $\mathfrak{F}_{\infty,\alpha}$ is separable. ■

Proposition 2.18. *Suppose $f\in\mathfrak{F}_{\infty,\alpha}, \alpha>0$. Then $f\in\mathfrak{F}^0_{\infty,\alpha}$ if and only if*

$$\lim_{r\to 1}\|f_r-f\|_{\mathfrak{F}_{\infty,\alpha}}=0, \tag{4}$$

where the power series expansion of $f(z)=\sum q^m a_m$ and so we can write

$$f_r(q)=f(rq)=\sum_{l=1}^{k}\sum_{|m|=l} r^{|m|}q^m a_m\ for\ all\ a_m\in\mathbb{H}^n,\ q\in\mathbb{B}^n\ and\ 0<r<1$$

with multi-index $m=(m_1,m_2,.....,m_n)$ and $|m|=|m_1|+|m_2|+.....+|m_n|$.

Proof. Let $I,J\in\mathbb{S}$ be such that $I\perp J$. Then for any $a_m\in\mathbb{H}^n$, there exist $a_{m,1}$ and $a_{m,2}$ in $\mathbb{C}_I^n$ such that $a_m=a_{m,1}+a_{m,2}J$. Now, $f\in\mathfrak{F}_{\infty,\alpha}$ implies $f\in\mathfrak{F}_{\infty,\alpha,I}$. So we can choose holomorphic functions f_1,f_2 in the complex space $\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}$ such that $Q_I[f]=f_1+f_2J$. Furthermore, $f(z)=\sum z^m a_{m,1}+\sum z^m a_{m,2}J=f_1(z)+f_2(z)J$. Thus, by Remark 2.15 and for $n=1,2$, we have

$$\begin{aligned}\|f_{r,n}-f_n\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}} &\leq \|f_r-f\|_{\mathfrak{F}_{\infty,\alpha,I}}\\ &\leq \|f_r-f\|_{\mathfrak{F}_{\infty,\alpha}}\\ &\leq 2\|f_r-f\|_{\mathfrak{F}_{\infty,\alpha,I}}\\ &\leq 2(\|f_{r,1}-f_1\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}}+\|f_{r,2}-f_2\|_{\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}}).\end{aligned}$$

Thus because of [28, Theorem 2], the condition (4) holds if and only if $f_1,f_2\in\mathfrak{F}^0_{\infty,\alpha,\mathbb{C}_I}$ which is equivalent to $f\in\mathfrak{F}^0_{\infty,\alpha}$. ■

Here, we investigate the t^{th}-derivative criterion for the space $f\in\mathfrak{F}^0_{\infty,\alpha}$ on a unit ball $\mathbb{B}$ in $\mathbb{H}$.

Definition 2.19. *For each $t\in\mathbb{N}\cup\{0\}$, we define the space $\mathfrak{F}^{(t)}$ of entire slice regular function f such that*

$$\lim_{|q|\to\infty}\frac{\left|\partial_{x_0}^{(t)}f(q)\right|}{(1+|q|)^t}e^{\frac{-\alpha}{2}|q|^2}=0,\ q\in\mathbb{B}.$$

Remark 2.20. Let $f\in SR(\mathbb{H},\mathbb{B})$ and $I,J\in\mathbb{S}$ be orthogonal imaginary units. Let f_1,f_2 be two holomorphic functions such that $Q_I[f]=f_1+f_2J$. Then for all $z\in\mathbb{B}_I$

$$|\partial_{x_0}^{(t)}f(q)|\leq|f_1^{(t)}(z)|+|f_2^{(t)}(z)|.$$

By Theorem 1.4, it follows that f lie in $\mathfrak{F}^{(t)}$ if and only if $f_l\in\mathfrak{F}^{(t)}_{\mathbb{C}_I}$ for $l=1,2$.

We can easily prove the following results.

Theorem 2.21. *Suppose* $0 \leq r < 1$, *if* $f \in \mathfrak{F}^{(t)}$, *then*

$$\lim_{r\to 1} \|f_r - f\|_{\mathfrak{F}_{\infty,\alpha}} = 0,$$

where $f_r(q) = f(rq) = \sum_{k=0}^{\infty} r^k q^k a_k$ *for all* $a_k \in \mathbb{H}$, $q \in \mathbb{B}$.

Proposition 2.22. $\mathfrak{F}^{(t)}$ *is the closure with respect to norm* $\|.\|_{\mathfrak{F}_{\infty,\alpha}}$ *of the set of quaternionic polynomials of the form*

$$P_k(q) = \sum_{m=0}^{k} r^m q^m a_m \text{ for all } a_m \in \mathbb{H},$$

for some positive integer k *and* $0 < r < 1$. *In particular,* $\mathfrak{F}^{(t)}$ *is separable.*

Theorem 2.23. *Let* $f \in SR(\mathbb{B}, \mathbb{H})$. *Then for any positive integer* t, $f \in \mathfrak{F}_{\infty,\alpha}$ *if and only if*

$$\frac{|\partial_{x_0}^{(t)} f(q)|}{(1+|q|)^t} e^{\frac{-\alpha}{2}|q|^2} \in l^{\infty}(\mathbb{H}). \tag{5}$$

Proof. Let $f \in \mathfrak{F}_{\infty,\alpha}$. Then $f \in \mathfrak{F}_{\infty,\alpha,I}$. Let $I, J \in$ with $I \perp J$. So, we can find holomorphic functions $f_1, f_2 : \mathbb{B} \cap \mathbb{C}_I \to \mathbb{C}_I$ such that $f_I = f_1 + f_2 J$. Now for any $z \in \mathbb{B}_I$, we have

$$\frac{|f_l^{(t)}(z)|}{(1+|z|)^t} e^{\frac{-\alpha}{2}|z|^2} \leq \frac{|\partial_{x_0}^{(t)} f(z)|}{(1+|z|)^t} e^{\frac{-\alpha}{2}|z|^2} \leq \frac{|f_1^{(t)}(z)|}{(1+|z|)^t} e^{\frac{-\alpha}{2}|z|^2} + \frac{|f_2^{(t)}(z)|}{(1+|z|)^t} e^{\frac{-\alpha}{2}|z|^2} \text{ for } l = 1, 2.$$

On applying [27, Theorem 1] to f_1 and f_2, it follows that the condition (5) holds if and only if $\frac{|f_l^{(t)}(z)|}{(1+|z|)^t} e^{\frac{-\alpha}{2}|z|^2} \in L^{\infty}(\mathbb{C}_I)$, $l = 1, 2$. Thus, we say that the above condition holds if and only if f_1, f_2 belong to complex space $\mathfrak{F}_{\infty,\alpha,\mathbb{C}_I}$ which is equivalent to $f \in \mathfrak{F}_{\infty,\alpha,I}$ and so $f \in \mathfrak{F}_{\infty,\alpha}$. ∎

References

[1] K. Abu-Ghanem, D. Alpay, F. Colombo, D. P. Kimsey, I. Sabadini, Boundary interpolation for slice hyperholomorphic Schur functions, *Integr. Equ. Oper. Theory,* **82** (2015), 223-248.

[2] K. Abu-Ghanem, D. Alpay, F. Colombo, I. Sabadini, Gleasons problem and Schur multipliers in the multivariable quaternionic setting, *J. Math. Anal. Appl.,* **425** (2015), 1083-1096.

[3] S. Adler, Quaterninoic Quantum Field Theory, *Oxford University Press*, 1995.

[4] D. Alpay, The Schur algorithm, reproducing kernel spaces and system theory, *Amer. Math. Soc.*, Providence, RI, 2001. Translated from the 1998 French original by Stephen S. Wilson, Panoramas et Synthèses.

[5] D. Alpay, V. Bolotnikov, F. Colombo, I. Sabadini, Self-mappings of the quaternionic unit ball: multiplier properties, Schwarz-Pick inequality, and Nevanlinna-Pick interpolation problem, **arXiv:1308.2658,** (2013).

[6] D. Alpay, V. Bolotnikov, F. Colombo, I. Sabadini, Interpolation problems for certain classes of slice hyperholomorphic functions, *Integr. Equ. Oper. Theory*, **86** (2016), 165-183.

[7] D. Alpay, F. Colombo, I. Sabadini, Schur functions and their realizations in the slice hyperholomorphic setting, *Integr. Equ. Oper. Theory,* **72** (2012), 253-289.

[8] D. Alpay, F. Colombo, I. Sabadini, Pontryagin-de Branges-Rovnyak spaces of slice hyperholomorphic functions, *J. Anal. Math.,* **121** (2013), 87-125.

[9] D. Alpay, F. Colombo, I. Sabadini, Inner product spaces and Krein spaces in the quaternionic setting in Recent Advances in Inverse Scattering, Schur Analysis and Stochastic Processes, Operator Theory: *Advances and Applications*, **244** (2015), 33-65.

[10] D. Alpay, F. Colombo, I. Sabadini, G. Salomon, Fock spaces in the slice hyperholomorphic setting, *In hypercomplex analysis: New Perspectives and Applications,* (2014), 43-59.

[11] D. Alpay, F. Colombo, I. Sabadini, T. Qian, The H^∞ functional calculus based on the S-spectrum for quaternionic operators and for n-tuples of noncommuting operators, **arXiv:1511.07629**, (2015).

[12] N. Arcozzi, G. Sarfatti, From Hankel operators to Carleson measures in a quaternionic variable, **arXiv:1407.8479,** (2014).

[13] N. Arcozzi, G. Sarfatti, Invariant metrics for the quaternionic Hardy space, *J. Geom. Anal.,* **25** (2015), 2028-2059.

[14] N. Arcozzi, G. Sarfatti, The orthogonal projection on slice functions on the quaternionic sphere, in Proceedings 30th International Colloquium on Group Theoretical Methods (F. Brackx, H. De Schepper and J. Vander Jeugt, editors), *Journal of Physics: Conference Series*, **597**.

[15] F. Colombo, J. O. Gonzalez-Cervantes, M.E. Luna-Elizarraras, I. Sabadini, M.V. Shapiro, On two approaches to the Bergman theory for slice regular functions, *Advances in Hypercomplex Analysis, Springer INdAM Series 1,* (2013), 39-54.

[16] F. Colombo, J. O. Gonzalez-Cervantes, I. Sabadini, On Slice biregular functions and isomorphisms of Bergman spaces, *Complex Var. Elliptic Equ.,* **57** (2012), 825-839.

[17] F. Colombo, J. O. Gonzalez-Cervantes, I. Sabadini, The C-property for slice regular functions and applications to the Bergman space, *Complex Var. Elliptic Equ.,* **58** (2013), 1355-1372.

[18] F. Colombo, I. Sabadini, D. C. Struppa, Noncommutative Functional Calculus, *Theory and Applications of Slice Regular Hyperholomorphic Functions,* Progress in Mathematics V. **289**, Birkhäuser Basel, 2011.

[19] F. Colombo, I. Sabadini, D. C. Struppa, Entire slice regular functions, **arXiv:1512.04215**, (2015).

[20] V. Fock, Verallgemeinerung und Lösung der Diracschen statistischen Gleichung, Zeitschrift für Physik **49** (1928), 339-357.

[21] G. Gentili, D. C. Struppa, A new approach to Cullen-regular functions of a quaternionic variable, *C. R. Math. Acad. Sci., Paris,* **342** (2006), 741-744.

[22] G. Gentili, D. C. Struppa, A new theory of regular functions of a quaternionic variable, *Adv. Math.,* **216** (2007), 279-301.

[23] S. Janson, J. Peetre and R. Rochberg, Hankel forms and the Fock spaces, *Revista Mat. Iberoamericana* **3** (1987), 61-138.

[24] S. Kumar, K. Manzoor, P. Singh, Composition operators in Bloch spaces of slice hyperholomorphic functions, *Adv. Appl. Clifford Algebras*, doi: 10.1007/s00006-016-0737-z, (2016).

[25] K. R. Parthasarathy, An Introduction to Quantum Stochastic Calculus, *Springer Science & Business Media*, 1992.

[26] G. Sarfatti, Elements of function theory in the unit ball of quaternions, Ph. D thesis, Universitá di Firenze, 2013.

[27] S. Ueki, Characterization for Fock-type spaces via higher order derivatives and its applications, *Complex Anal. Oper. Theory,* **8** (2014), 1475-1486.

[28] S. Ueki, Higher order derivative characterization for Fock-type spaces, *Integr. Equ. Oper. Theory,* **84** (2016), 89-104.

[29] C. M. P. C. Villalba, F. Colombo, J. Gantner, J. O. G. Cervantes, Bloch, Besov asnd Dirichlet spaces of slice hyerhlomorphic functions, *Complex Anal. Oper. Theory,* **9** (2015), 479-517.

[30] R. Wallsten, The S^p-criterion for Hankel forms on the Fock space, $0 < p < 1$, *Math. Scand.,* **64** (1989), 123-132.

[31] K. Zhu, Operator Theory in Function Spaces, Marcel Dekker, New York, 1990.

[32] K. Zhu, Analysis on Fock Spaces, *Graduate Texts in Mathematics* **263**, Springer, New York, 2012.

Department of Mathematics, Central University of Jammu, Jammu 180 011, INDIA.
E-mail address: sanjaykmath@gmail.com

Department of Mathematics, Central University of Jammu, Jammu 180 011, INDIA.
E-mail address: khalidcuj14@gmail.com